



Still not right after a concussion? Check your vision.

A concussion can disrupt how the brain aims, focuses and moves the eyes, even when scans and the eye chart are normal. These problems are common, measurable and treatable, and starting early helps.

Check anything you've noticed since the injury

- | | |
|---|---|
| ☐ Reading makes symptoms worse | ☐ Losing your place or rereading lines |
| ☐ Headaches with screens or reading | ☐ Words blur, move or double |
| ☐ Ordinary light is uncomfortable | ☐ Queasy when scrolling or watching motion |
| ☐ Dizzy in stores, crowds or traffic | ☐ Trouble judging distance |
| ☐ Hard to concentrate on a page | ☐ Exhausted by the end of the day |

Don't wait for everything else to clear

Once your doctor has checked you, visual symptoms can be measured right away. Injuries from years ago still count. Your care team gets the results, with your permission.



Book an evaluation
visionlearncenter.com
(561) 462-1245

Go to an emergency room for worsening headache, repeated vomiting, confusion, weakness or sudden double vision.

Referral to Vision & Learning Center

CUT HERE AND BRING TO YOUR VISIT

Patient's name

Date of birth / date of injury

Referring provider

Practice and phone

Findings: ☐ Receded near point of convergence ☐ Symptoms on ocular motor screen
☐ Visual-vestibular ☐ Stalled return to learn / work / play

Offices across South Florida · Children and adults